

Information for staff on Pneumocystis Pneumonia (PcP): Outbreak prevention and management

The purpose of this information is to provide an aide-memoire for clinical staff and infection prevention and control teams (IPCT), who may be involved in the outbreak prevention and management of PcP.

All staff should be familiar with [National Infection Prevention and Control Manual](#) (NIPCM), standard infection control precautions (SICPs) and transmission based precautions (TBPs).

The following advice is supplementary and provides details of specific actions necessary to prevent and manage PcP outbreaks. It should be noted that there is currently no up-to-date national or European guidance relating to the prevention and management of PcP in high-risk patients.

Outbreak prevention

What is PcP?

PcP is caused by the opportunistic fungus *Pneumocystis jirovecii* (formerly *P. carinii*) that affects the respiratory system. Typical presentation includes fever, cough and shortness of breath¹, which develop after a relatively long incubation period, ranging from approximately three to 12 weeks.^{2;3} Transmission is considered to occur via the airborne route, although there is a lack of consensus regarding the exact mechanism.⁴

Which patient groups are considered high-risk for infection?

Severely immunocompromised patients including those with HIV (CD4+ counts < 200 cells/μl), haematological and solid malignancies receiving cytotoxic chemotherapies, solid organ transplant patients, those on high dose steroids and patients treated with immunosuppressive regimens for inflammatory conditions.^{5;6} Since PcP is unlikely to affect immunocompetent hosts, consider excluding causes of underlying immunosuppression in those diagnosed with the infection.⁶

What Transmission Based Precautions should be applied?

P. jirovecii is considered to mainly pose an infection risk to immunosuppressed patients, therefore apply the following precautions to prevent ongoing transmission in this patient group.

Optimal patient placement

- Isolate patients with PcP in a single ensuite room in high-risk units or settings, (for example ICU, PICU, NICU, oncology, haematology, renal), until resolution of symptoms or discharge from hospital. ^{4,7}
- Avoid placing patients diagnosed with PcP in the same ward or healthcare area as immunocompromised patients.⁸

PPE to be worn by healthcare staff

- Transmission risk to immunocompromised patients is via the droplet route due to symptoms associated with PcP, for example cough. There is currently insufficient evidence of PcP healthcare staff acquisition from colonised patients ⁹⁻¹¹ Staff must wear PPE appropriate to the risk associated with droplet transmission.

Mask use by patients

- In an outbreak, all immunocompromised patients should be encouraged to wear single-use, fluid resistant masks during transport between wards or clinical areas to reduce the likelihood of cross-transmission.^{4,7}

How can PcP be prevented?

Use of prophylaxis is considered to be the most successful prevention strategy for PcP. Prophylaxis with co-trimoxazole (as a first line agent) of appropriate at-risk patient groups is recommended. Dapsone or pentamidine are typically prescribed as alternatives.^{1;4-6} The duration of prophylaxis varies depending on the patient's condition and level of immunosuppression.^{1;6}

How should PcP be managed?

Successful management of infection is attributed to appropriate treatment of affected patients. As with prophylaxis, patients should be treated with co-trimoxazole as the first-line agent (a number of alternatives are also available).^{4;6} Treatment duration is approximately two to three weeks.^{5;6} In addition, it may be necessary to consider prophylaxis for all immunosuppressed contacts,⁴ including patients who have been discharged, up to 12 weeks prior to recognition of the outbreak (as determined by the incubation period).

Outbreak Management

How is a potential PcP outbreak identified?

A single healthcare-associated case in a high-risk patient will require investigation and should be considered as part of 'alert organism' infection prevention and control teams (IPCT) actions.^{4,7} An outbreak may be suspected if the incidence of infection is higher than normally expected in a given healthcare area over a specified time period, and where two or more cases may have a potential link in time and place.⁷

Outbreaks caused by *P. jirovecii* are often recognised late. This is generally considered to be due to the long incubation period associated with the infection, which is typically followed by a non-specific presentation and often initially mistaken for other infectious causes.⁷

In line with [Chapter 3](#) of the NIPCM following detection or recognition of an incident, a member of IPCT or health protection team (HPT) should undertake an outbreak assessment and investigation.⁷

Clinical management of contacts

In the absence of national or European Guidance relating to the Prevention and Management of PcP in high-risk patients, consider the following:

- exposed contacts should be clinically assessed for risk
- offer prophylaxis outlined as above, or
- for at-risk patients that maybe also have a higher risk of toxicity related to prophylactic treatment, implement close monitoring through screening of the contacts' respiratory samples and provide early treatment if positive patients are identified.⁴

Investigation of sources of transmission

During an outbreak it is prudent to consider a hypothesis that will inform an investigatory review of potential sources. Consider the following where ongoing transmission is identified:

- screening – to identify any asymptomatic carriers, and
- [ventilation system-associated](#) risk.

Reference List

- (1) Morris A, Norris K. Colonization by *Pneumocystis jirovecii* and Its Role in Disease. *Clinical Microbiology Reviews* 2012;297-317.
- (2) Schmolt S, Schuегger R, Wendler T, Huber I, Sollner H, Hogardt M, et al. Molecular evidence of nosocomial *Pneumocystis jirovecii* transmission among 16 patients after kidney transplantation. *Journal of Clinical Microbiology* 2008;46(3):966-71.
- (3) Le GS, Damiani C, Rouille A, Grall A, Treguer L, Virmaux M, et al. A cluster of *Pneumocystis* infections among renal transplant recipients: molecular evidence of colonized patients as potential infectious sources of *Pneumocystis jirovecii*. *Clinical Infectious Diseases* 2012 Apr;54(7):e62-e71.
- (4) Yiannakis EP, Boswell TC. Systematic review of outbreaks of *Pneumocystis jirovecii* pneumonia: Evidence that *P. jirovecii* is a transmissible organism and the implications for healthcare infection control. *Journal of Hospital Infection* 2016;93(1):01.
- (5) BMJ Best Practice. *Pneumocystis jirovecii* pneumonia. 2016.
- (6) American Thoracic Society. An Official American Thoracic Society Statement: Treatment of Fungal Infections in Adult Pulmonary and Critical Care Patients. *American Journal of Respiratory Medicine* 2011;183:96-128.
- (7) Health Protection Scotland. [National Infection Prevention and Control Manual. Chapter 3: Healthcare Infection Incidents, Outbreaks and Data Exceedance](#). 2019
- (8) CDC. Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings. 2007.
- (9) Mori S, Sugimoto M. *Pneumocystis jirovecii* infection: An emerging threat to patients with rheumatoid arthritis. *Rheumatology (United Kingdom)* 2012;51(12):kes244.
- (10) Mori S, Cho I, Sugimoto M. A cluster of *Pneumocystis jirovecii* infection among outpatients with Rheumatoid Arthritis. *The Journal of Rheumatology* 2010;37(7):1547-8.
- (11) Phipps LM, Chen SC, Kable K, Halliday CL, Firacative C, Meyer W, et al. Nosocomial *Pneumocystis jirovecii* pneumonia: lessons from a cluster in kidney transplant recipients. *Transplantation* 2011 Dec 27;92(12):1327-34.
- (12) Centre for Disease Control. [Pneumocystis Pneumonia Basics](#). 2024